

Systems of Nonlinear Eqs.

A. DONEV, Spring 2021

Consider now a system of two nonlinear eqs. in 2 variables:

$$\begin{cases} f_1(x_1, x_2) = 0 \\ f_2(x_1, x_2) = 0 \end{cases}$$

I imagine we are given a guess for the solution $x_1^{(k)}$ and $x_2^{(k)}$ and we want to improve it to get $x_{1,2}^{(k+1)}$ (Worksheet 2)

There is no equivalent of bisection in higher dimensions!

(1)

1D is very special because the real line is ordered! But the plane (or $\mathbb{R}^{d>1}$) is not ordered.

Newton's method, however, generalizes easily since it is based on a Taylor series!

$$f_1(x_1, x_2) = f_1(x_1^{(k)}, x_2^{(k)}) + \frac{\partial f_1}{\partial x_1}(x_1^{(k+1)} - x_1^{(k)}) + \frac{\partial f_1}{\partial x_2}(x_2^{(k+1)} - x_2^{(k)})$$

and similarly for f_2

(2)

Write this using matrix notation as

$$\vec{f} = (f_1, f_2)^T$$

$$\vec{x} = (x_1, x_2)^T$$

$$\vec{f}(\vec{x}) = \vec{0}$$

$$f(\vec{x}^{(k+1)}) = f(\vec{x}^{(k)}) +$$

$$\begin{bmatrix} \partial f_1 / \partial x_1 & \partial f_1 / \partial x_2 \\ \partial f_2 / \partial x_1 & \partial f_2 / \partial x_2 \end{bmatrix} \begin{bmatrix} x_1^{(k+1)} - x_1^{(k)} \\ x_2^{(k+1)} - x_2^{(k)} \end{bmatrix}$$

$$\vec{f}(\vec{x}^{(k+1)}) = \vec{f}(\vec{x}^{(k)}) + \vec{J}(\vec{x}^{(k)}) (\vec{x}^{(k+1)} - \vec{x}^{(k)})$$

$$J_{ij} = \frac{\partial f_i}{\partial x_j}$$

Jacobian matrix

(3)

Idea in Newton's method:

Replace $f(\vec{x})$ by its first-order Taylor series and find root of that linear function:

$$f(\vec{x}^{(k+1)}) = \vec{0}$$

$$J(\vec{x}^{(k)}) \Delta X^{(k)} = -f(\vec{x}^{(k)})$$

Newton's method

$$\vec{x}^{(k+1)} = \vec{x}^{(k)} + \Delta X^{(k)}$$

Linear system to be solved

each iteration using LU factorization. Books may write

$$\vec{x}^{(k+1)} = \vec{x}^{(k)} - (J^{(k)})^{-1} f^{(k)} \text{ but}$$

use backslash instead

(4)

When to stop?

Termination or stopping criterion

Two options:

① Stop if $\|f^k\| < \epsilon$

\uparrow
1, 2, ∞
tolerance

② Stop if $\|\Delta x^k\| < \epsilon$

\downarrow
1, 2, ∞

Options are related but not the same

Option 2: $\|(\mathcal{J}^k)^{-1} f^k\| < \epsilon$

estimate $\|(\mathcal{J}^k)^{-1}\| \|f^k\| < \epsilon$

$\|f^k\| < \epsilon / \|(\mathcal{J}^k)^{-1}\|$ (4 1/2)

Note: Linear systems are at the heart/core of NA!

For scalar functions of many variables

$$f: \mathbb{R}^n \rightarrow \mathbb{R}$$

Taylor series to second order is

$$f(\vec{x} + \Delta\vec{x}) = f(\vec{x}) + \vec{g}^T \Delta\vec{x} + \frac{1}{2} (\Delta\vec{x})^T \mathbf{H} \Delta\vec{x} + O(\|\Delta\vec{x}\|^3)$$

where gradient (transpose of Jacobian) is column vector

$$\vec{g} = \nabla_{\vec{x}} f = \vec{\nabla} f = \left[\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \right]^T$$

(5)

and Hessian matrix is

$$\mathbb{H} = \nabla_x^2 f$$

$$H_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j} = H_{ji}$$

if f is twice continuously differentiable.

This is one of the reasons symmetric matrices are important!

For $f: \mathbb{R}^m \rightarrow \mathbb{R}$

gradient (vector) replaced by Jacobian (matrix)

and

Hessian (matrix) replaced by rank-3 tensor

(6)

Convergence of Newton method

$$\text{Let } \vec{f}(\vec{x}) = \vec{0}$$

\uparrow
root

$$\vec{e}^{(k)} = \vec{x}^{(k)} - \vec{x} = \text{error at step } k$$

Near the root

$$\vec{J}(\vec{x}^{(k)}) \approx \vec{J}(\vec{x})$$

So evaluate derivatives at root
in analysis

$$\vec{f}(\vec{x}^{(k)}) = \vec{f}(\vec{x}) + \vec{J}(\vec{x}) \vec{e}^{(k)} + \frac{1}{2} (\vec{e}^{(k)})^T \vec{H}(\vec{e}^{(k)})$$

$$\begin{aligned} \vec{e}^{(k+1)} &= \vec{x}^{(k+1)} - \vec{x} = \vec{J}^{-1} \vec{f}(\vec{x}^{(k)}) \\ &= \left(\vec{x}^{(k)} - \vec{x} \right) - \vec{J}^{-1} \vec{f}(\vec{x}^{(k)}) \end{aligned} \quad (7)$$

$$\begin{aligned} \Rightarrow \vec{e}^{(k+1)} &= \vec{e}^{(k)} - \mathbb{J}^{-1} \vec{f}(\vec{x}^{(k)}) \\ &= \vec{e}^{(k)} - \mathbb{J}^{-1} \left[\vec{f}(\vec{x}^{(k)}) + \mathbb{J} \vec{e}^{(k)} + \frac{1}{2} (\vec{e}^{(k)})^T \mathbb{H} (\vec{e}^{(k)}) \right] \end{aligned}$$

$$\Rightarrow \vec{e}^{(k+1)} = -\frac{1}{2} (\mathbb{J}^{-1}) (\vec{e}^{(k)})^T \mathbb{H} (\vec{e}^{(k)})$$

rank 3
tensor

$$\|\vec{e}^{(k+1)}\| \leq \frac{\|\mathbb{J}^{-1}\| \|\mathbb{H}\|}{2} \|\vec{e}^{(k)}\|^2$$

quadratic convergence

as long as Jacobian is not (nearly) singular

⑧

Compare this to 1D
analysis

$$e^{(k+1)} \approx -\frac{1}{2} \frac{f''(x)}{f'(x)} (e^{(k)})^2$$

So if Newton's method
in higher dims converges,
it will eventually converge
fast once it gets
sufficiently close to the root.

But Newton's method
requires a good guess to
converge (and since no
bisection there is no easy
way to get a good initial
guess).

⑨